Renal Involvement and Early Prognosis in Patients with COVID-19 Pneumonia

Guangchang Pei,¹ Zhiguo Zhang ^(b),^{2,3} Jing Peng,⁴ Liu Liu,¹ Chunxiu Zhang,¹ Chong Yu,¹ Zufu Ma,¹ Yi Huang,¹ Wei Liu,¹ Ying Yao,¹ Rui Zeng,¹ and Gang Xu¹

¹Division of Nephrology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

²School of Medicine and Health Management, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

³Hubei Provincial Research Center for Health Technology Assessment, Wuhan, China

⁴Department of Laboratory Medicine, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China

ABSTRACT

Background Some patients with COVID-19 pneumonia also present with kidney injury, and autopsy findings of patients who died from the illness sometimes show renal damage. However, little is known about the clinical characteristics of kidney-related complications, including hematuria, proteinuria, and AKI.

Methods In this retrospective, single-center study in China, we analyzed data from electronic medical records of 333 hospitalized patients with COVID-19 pneumonia, including information about clinical, laboratory, radiologic, and other characteristics, as well as information about renal outcomes.

Results We found that 251 of the 333 patients (75.4%) had abnormal urine dipstick tests or AKI. Of 198 patients with renal involvement for the median duration of 12 days, 118 (59.6%) experienced remission of pneumonia during this period, and 111 of 162 (68.5%) patients experienced remission of proteinuria. Among 35 patients who developed AKI (with AKI identified by criteria expanded somewhat beyond the 2012 Kidney Disease: Improving Global Outcomes definition), 16 (45.7%) experienced complete recovery of kidney function. We suspect that most AKI cases were intrinsic AKI. Patients with renal involvement had higher overall mortality compared with those without renal involvement (28 of 251 [11.2%] versus one of 82 [1.2%], respectively). Stepwise multivariate binary logistic regression analyses showed that severity of pneumonia was the risk factor most commonly associated with lower odds of proteinuric or hematuric remission and recovery from AKI.

Conclusions Renal abnormalities occurred in the majority of patients with COVID-19 pneumonia. Although proteinuria, hematuria, and AKI often resolved in such patients within 3 weeks after the onset of symptoms, renal complications in COVID-19 were associated with higher mortality.

JASN 31: •••-, 2020. doi: https://doi.org/10.1681/ASN.2020030276

The outbreak of coronavirus disease has rapidly evolved into a global pandemic.^{1–7} This novel coronavirus is named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2).⁸ The World Health Organization (WHO) officially named the disease caused by SARS-CoV-2 as coronavirus disease 2019 (COVID-19). The initial clinical sign for the detection of COVID-19 was pneumonia.³ However, other organ damages were also reported.^{6,9} Some cases of COVID-19 pneumonia presented with kidney injury,⁴ and pathologic findings from autopsies also revealed renal damage from the corpses of patients with COVID-19¹⁰; thus, SARS-CoV-2 may include kidney tropism. In this study, we reported renal involvement and their early prognosis of patients with COVID-19 pneumonia admitted to Tongji Hospital in Wuhan, China, in terms of proteinuria, hematuria, and AKI.

METHODS

Study Design and Participants

For this retrospective, single-center study, we included adult patients from January 28 to February 9, 2020, at Tongji

Received March 9, 2020. Accepted April 12, 2020.

G.P. and Z.Z. contributed equally to this work.

Published online ahead of print. Publication date available at www.jasn.org.

Correspondence: Dr. Ying Yao or Dr. Rui Zeng, Division of Nephrology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, No. 1095, Jiefang Avenue, Wuhan 430030, China. E-mail: yaoyingkk@126.com or zengr126@126.com

Copyright $\ensuremath{\mathbb{C}}$ 2020 by the American Society of Nephrology

Hospital of Sino-French New City District, which was one of the designated hospitals solely for the treatment of patients with COVID-19 in Wuhan, China. The final follow-up for this study ended on February 23, 2020. All patients who were diagnosed as having COVID-19 pneumonia according to the Diagnosis and Treatment Protocols of Pneumonia caused by Novel Coronavirus (SARS-CoV-2) by the National Health Commission of China (Trial Version 7)¹⁰ were screened. Specifically, the diagnosis criteria were as follows: (1) fever or respiratory symptoms, (2) leukopenia or lymphopenia, and (3) computed tomography (CT) scan showing radiographic abnormalities in lung. Those with two or more diagnosis criteria (lung involvement is necessary) and a positive result on high-throughput sequencing or RT-PCR assay were diagnosed as having COVID-19 pneumonia.¹⁰ Exclusion criteria were as follows: (1) patients with CKD or suspected CKD, with at least one abnormal urine test including proteinuria and hematuria in the 3 months before admission; (2) patients with maintenance dialysis, renal transplantation, or peak serum creatinine of $<53 \ \mu mol/L;$ (3) patients with time from onset of disease to admission date of >14 days, given that most patients achieved radiologic remission after 14 days from onset.11 The study was approved by Tongji Hospital Ethics Committee of Huazhong University of Science and Technology (approval number TJ-C20200132). Informed consent was waived in the light of the urgency of data collection.

Data Collection and Measurements

We obtained demographic, epidemiologic, clinical, laboratory, and radiologic characteristics, and treatment and renal outcome data from electronic medical records. The data were reviewed by two physicians (G.P. and R.Z.). The onset date was defined as the day when the symptom was noticed. As of February 9, 2020, a total of 467 patients with COVID-19 were admitted and screened. A total of 333 patients with urine dipstick test on the first morning after

admission, or with AKI on admission and during the hospital stay, were included in this study. Among the 333 patients, 82 patients without proteinuria or hematuria on admission and without AKI were excluded. Further, 53 patients without serial monitoring of urine dipstick tests were excluded. Finally, 198 patients with serial monitoring of urine dipstick tests, or with AKI, were included in the cohort (Supplemental Figure 1). It should be noted that 19 of the 35 patients with AKI and 235 of the 298 patients without AKI have been described previously by Cheng et al.12

Patients with possible COVID-19 pneumonia were admitted and guarantined, and throat swab samples were collected and detected in the hospital using a quantitative real-time RT-PCR to confirm SARS-CoV-2, as follows: A Viral Nucleic Acid Kit (Tianlong Science & Technology Co., Ltd., Xi'An, China) was used to extract nucleic acids from throat swab samples and a SARS-CoV-2 detection kit (DAAN GENE Co., Guangzhou, China) was used to detect the ORF1ab gene (nCovORF1ab) and the N gene (nCoV-NP) according to the manufacturer's instructions, using real-time RT-PCR. If the circulation threshold (Ct) values for both genes were >40 or had no typical amplification curves, and the internal control performed well, it was considered that the SARS-CoV-2 RNA was not present. If the Ct values for both genes were ≤ 40 , the gene detection results were considered as positive. If only one gene had a Ct value of ≤ 40 , and the other one had no typical amplification curve, a repeat experiment was performed, with consistent results considered as positive for SARS-CoV-2 RNA.

All patients were given a chest CT scan before or after admission to hospital, and a repeat chest CT scan was obtained at 5–10 day intervals. Laboratory data consisted of complete blood counts, liver and renal function, coagulation function, high-sensitivity C-reactive protein (CRP), erythrocyte sedimentation rate, and serum cytokines. Serial monitoring of these laboratory tests

Significance Statement

Some cases of COVID-19 pneumonia have presented with kidney injury, and autopsy findings for patients with COVID-19 have revealed renal involvement. In this retrospective, single-center study of Chinese patients with COVID-19 pneumonia, 251 of 333 patients (75.4%) presented with renal complications, including proteinuria, hematuria, and AKI. After the median duration of 12 days of follow-up, nearly half of patients with AKI recovered from AKI within 3 weeks of onset of infection. However, patients with renal involvement had higher overall mortality (11.2%) compared with patients without renal involvement (1.2%). Adverse short-term outcomes of renal involvement were associated with severity of COVID-19 pneumonia. These findings indicate that although early renal abnormalities often resolve in such patients, intensive support and careful monitoring of severe or critical illness is appropriate for COVID-19 pneumonia with renal complications.

was performed for each patient according to the patient's clinical progress.

Definitions

COVID-19 pneumonia was classified into four types, namely, mild, moderate, severe, and critically ill, according to Diagnosis and Treatment Protocols of Pneumonia caused by Novel Coronavirus (SARS-CoV-2) by the National Health Commission of China (Trial Version 7).10 Moderate COVID-19 pneumonia was defined as fever, respiratory syndrome, and radiologic lung findings.¹⁰ Severe COVID-19 pneumonia was defined as meeting any of the following conditions: (1) respiratory rate ≥ 30 breaths/min, (2) oxygen saturation \leq 93% in a resting state, (3) Arterial oxygen partial pressure(PaO2) / Fractional inspired oxygen(FiO2) ratio \leq 300 mm Hg, and (4) a 50% increase in chest radiologic abnormalities in 24-48 hours.^{10,12} Critically ill COVID-19 pneumonia was defined as either (1) respiratory failure need for mechanical ventilation, (2) shock, or (3) organ failure need for ICU admission.10,12

For the diagnosis of AKI, we referred to the 2012 Kidney Disease: Improving Global Outcomes (KDIGO) definition and the expanded criteria used by Yang et al.13,14 First, we screened patients with suspected AKI using 2012 KDIGO definition as the major screening criteria. However, patients without previous baseline serum creatinine and those who did not meet the 2012 KDIGO criteria at admission but their increased serum creatinine was 1.5 times above the baseline with intervals longer than 7 days, would be excluded according to the 2012 KDIGO definition, which might potentially underestimate overall occurrence of AKI. For those who had no repeated serum creatinine within 7 days or with recovering AKI, we expanded the screening criteria to an increase or decrease in serum creatinine by \geq 0.3 mg/dl during hospital stay. Second, these suspected AKI were reviewed on a case-by-case basis to confirm the diagnosis. The identification criteria included the 2012 KDIGO definition of AKI (KDIGO criteria). For those who had no repeated serum creatinine within 7 days or with recovering AKI, we expanded the identification criteria of AKI to an increase or decrease in serum creatinine by 50% during hospital stay (using serum creatinine concentration at admission as a baseline; we looked for increases or decreases in creatinine relative to this value), with or without oliguria. We defined renal recovery as serum creatinine decreasing to below threshold or to the baseline. The staging of AKI was on the basis of 2012 KDIGO criteria.14

Patients were considered to have prerenal AKI when the rise in serum creatinine concentration had been caused by low renal perfusion, and creatinine recovered rapidly to baseline after volume administration within 3 days.¹⁵ Rhabdomyolysis was diagnosed on the basis of medical history, and elevated serum creatinine phosphokinase (CK) levels more than five times the upper limit of normal and/or serum myoglobin >150 ng/ ml.16,17 The elevations in CK were not caused by myocardial infarction, cerebral vascular disease, or neuromuscular disease. As no obstructive AKI was diagnosed, we classified the other AKI in this study as suspected intrinsic AKI, which

might include acute tubular necrosis (ATN) and nephrotoxic AKI.

Patients with CKD were excluded. CKD was defined as eGFR<60 ml/min per 1.73 m² or urine microalbumincreatinine ratio \geq 30 µg/mg at least 3 months before admission.¹⁸ If there was only one outpatient GFR <60 ml/ min per 1.73 m², we evaluated CKDrelated complications such as anemia and mineral and bone disorders. The patients who had CKD-related complications were considered as having suspected CKD and were also excluded.

Proteinuria and hematuria were defined as more than trace albumin or blood on urine dipstick tests, which were collected and detected on the first morning after admission and during the observation period. Remission of proteinuria and hematuria was defined as protein and blood were negative on urine dipsticks. Recovery of AKI was defined as complete recovery of kidney function.

Statistical Analyses

Data conforming to normal distribution were presented as mean±SD, or median and quartiles for non-normal distribution. Rate comparisons were performed by chi-squared test. t Test, Wilcoxon rank-sum test, Wilcoxon signed-rank test, or Kruskal-Wallis test were used to compare means across groups according to the number of group and distribution of variable. To estimate the degree of correlation between proteinuria or hematuria and clinical variables, Spearman rank correlation coefficient was used. Stepwise multivariate binary logistic regression was used to select and estimate the association between proteinuria, hematuria, or AKI remission and variables that were clinically relevant on grounds of professional knowledge and were statistically significant in preliminary univariate binary logistic regression. Specifically, variables with P < 0.05 in the univariate analysis were entered into multivariate analysis to select the predictors (inclusion criterion was P<0.05 and exclusion criterion was P < 0.10). Results are presented as odds ratios (ORs) with 95% confidence

intervals (95% CIs) and *P* values. All statistical analyses were performed using SPSS version 20.0 software (IBM Corp.).

RESULTS

Baseline Characteristics

A total of 333 patients were included in the study. The mean of age was 56.3 years (Table 1), and 54.7% (182 of 333) patients were men. The median duration from onset to admission was 9 (7–11) days (Table 1). The prevalence of hypertension and diabetes was 32.2% (107 of 333) and 22.9% (76 of 333), respectively. Compared with moderate cases, patients with severe or critically ill COVID-19 pneumonia were older, more likely to experience dyspnea, and more likely to have hypertension, diabetes, and angiotensinconverting enzyme inhibitor/angiotensin receptor blocker (ACEI/ARB) treatment history (Table 1).

Proteinuria, Hematuria, and AKI

On admission, of the 333 patients, 75.4% (251 of 333) patients had renal involvement, 65.8% (219 of 333) patients presented with proteinuria, and 41.7% (139 of 333) patients had hematuria (Table 1). The incidence of AKI in the overall cohort was 4.7% (22 of 467) by KIDGO criteria and 7.5% (35 of 467) by expanded criteria. Greater incidence of proteinuria (81.2% and 85.7%, respectively, versus 43.8%) and hematuria (39.1% and 69.6%, respectively, versus 33.3%) were demonstrated in patients with severe or critically ill COVID-19 pneumonia (Table 1). Among the 333 patients, the patients with AKI had higher incidence rate of proteinuria (88.6% versus 63.1%) and hematuria (60% versus 41.7%) compared with the non-AKI group (Supplemental Table 1). A total of 42.9% (24 of 56) critically ill cases developed AKI during the hospital stay (Table 1).

The number of patients with AKI was 22 by KDIGO criteria and 35 by expanded criteria (Table 2). According to KDIGO criteria, AKI occurred after admission in 19 (86.4%) of 22 patients with AKI. Rhabdomyolysis-induced

Table 1.	Clinical characteristics	of p	oatients	with	COVID-19
----------	--------------------------	------	----------	------	----------

Variables	All Patients	Moderate	Severe	Critically III	P Value
N	333	144 (43.2%)	133 (39.9%)	56 (16.8%)	
Days from onset	9 (7–11)	8 (6–11)	10 (7–11)	8 (6–11)	0.12ª
Age, yr	56.3±13.4	50.9±12.5	58.1±12.3	63.1±11.0	<0.001 ^a
Male patient, %	182/333 (54.7%)	67/144 (46.5%)	79/133 (59.4%)	36/56 (64.3%)	0.03 ^b
Fever, %	301/333 (90.4%)	126/144 (87.5%)	123/133 (92.5%)	52/56 (92.9%)	0.16 ^b
Cough, %	229/333 (68.8%)	98/144 (68.1%)	94/133 (70.7%)	37/56 (66.1%)	0.94 ^b
Dyspnea, %	190/333 (57.1%)	71/144 (49.3%)	85/133 (63.9%)	34/56 (60.7%)	0.05 ^b
Diarrhea, %	108/333 (32.4%)	42/144 (29.2%)	41/133 (30.8%)	25/55 (49.1%)	0.06 ^b
Hypertension, %	107/332 (32.2%)	37/144 (25.7%)	43/133 (32.3%)	27/55 (49.1%)	0.003 ^b
ACEI/ARB history, %	37/321 (11.5%)	12/143 (8.4%)	14/126 (11.1%)	11/52 (21.2%)	0.02 ^b
ACEI/ARB	11/26	0/12	6/8	5/6	
Diabetes, %	76/332 (22.9%)	20/144 (13.9%)	32/133 (24.1%)	24/55 (43.6%)	<0.001 ^b
SPO ₂ (%)	93 (91.0–97.0)	97.0 (96.0–98.0)	92.0 (90.0–93.0)	89.5 (80.0–93.0)	<0.001ª
Systolic BP, mm Hg	126 (115–138)	124 (114–132)	128 (116–141)	135 (118–145)	0.001ª
Diastolic BP, mm Hg	78 (72–86)	77 (72–86)	78 (71–87)	79 (73–86)	0.76ª
Blood sugar, mmol/L	6.6 (5.7-8.1)	6.0 (5.4–7.1)	6.9 (5.9–8.6)	7.8 (6.3–12.4)	<0.001ª
CRP, mg/L	44.1 (16.3–90.8)	22.1 (7.5-45.3)	58.1 (32.5–101.4)	69.7 (42.9–119.6)	<0.001ª
Erythrocyte sedimentation rate, mm/h	31.0 (18.0–58.8)	27.0 (14.3–50.3)	39.0 (22.0-64.0)	31.0 (22.0–58.0)	0.004ª
ALT, U/L	24.0 (15.0–38.3)	20.0 (13.0–33.0)	26.0 (18.5–39.5)	29.0 (18.0-43.0)	0.001ª
AST, U/L	31.0 (22.0–48.0)	24.0 (20.0–38.0)	33.0 (24.0–50.5)	40.5 (25.5–62.3)	<0.001ª
HsTnI, pg/ml	5.4 (3.3–14.0)	4.6 (2.5–7.0)	4.8 (3.1–11.7)	11.4 (6.3–21.6)	<0.001 ^a
NT-proBNP, pg/ml	139.0 (52.0–392.5)	76.0 (23.0–152.0)	157.5 (55.5–333.8)	372.5 (155.3–758.3)	0.002 ^a
Serum albumin, g/L	34.5±4.7	36.6±4.9	33.6±4.2	32.9±4.3	<0.001ª
BUN, mmol/L	4.3 (3.2–5.7)	3.9 (3.1–5.0)	4.4 (3.2–5.5)	5.9 (4.6–8.6)	<0.001ª
SCR, μ mol/L	70.0 (57.0–84.0)	66.5 (56.0–81.0)	69.0 (57.0–84.0)	77.0 (60.0–89.0)	<0.001ª
Prothrombin time, sec	14.1 (13.5–14.8)	14.0 (13.2–14.5)	14.3 (13.5–14.9)	14.4 (13.8–14.9)	0.004ª
D-dimer, mg/L	0.73 (0.40–1.33)	0.47 (0.32-1.10)	0.80 (0.57-1.33)	1.25 (0.64–5.90)	<0.001ª
Neutrophils, 10 ⁹ /L	3.79 (2.57–5.45)	2.99 (2.14–4.13)	4.06 (2.84–5.56)	5.79 (3.79–8.59)	<0.001ª
Lymphocytes, 10 ⁹ /L	0.86 (0.63–1.20)	1.01 (0.72–1.44)	0.78 (0.60–1.05)	0.55 (0.43–0.80)	<0.001ª
Eosinophils, 10 ⁹ /L	0.00 (0.00-0.01)	0.00 (0.00-0.09)	0.00 (0.00-0.06)	0.00 (0.00-0.01)	<0.001ª
Monocytes, 10 ⁹ /L	0.39 (0.28–0.51)	0.41 (0.30–0.53)	0.36 (0.26–0.52)	0.38 (0.22–0.49)	0.16ª
Serum $TNF\alpha$, pg/ml	8.6 (7.0–10.7)	8.1 (6.8–9.8)	9.3 (7.6–11.6)	9.3 (6.8–11.1)	0.09ª
Serum IL-10, pg/ml	5.7 (5.0–10.2)	5.0 (5.0-8.1)	6.4 (5.0–10.4)	8.0 (5.0–15.1)	0.01ª
Serum IL-6, pg/ml	19.9 (8.0-45.4)	13.2 (3.8–23.1)	27.1 (11.8–60.0)	32.8 (17.8–62.6)	<0.001ª
Serum IL-2R, U/ml	669 (459–963)	546 (455–743)	766 (595–1050)	1026 (378–1260)	<0.001ª
Proteinuria, %	219/333 (65.8%)	63/144 (43.8%)	108/133 (81.2%)	48/56 (85.7%)	<0.001 ^b
Hematuria, %	139/333 (41.7%)	48/144 (33.3%)	52/133 (39.1%)	39/56 (69.6%)	<0.001 ^b
AKI, %	35/333 (10.5%)	5/144 (3.5%)	6/133 (4.5%)	24/56 (42.9%)	<0.001 ^b
Renal involvement, %	251/333 (75.4%)	89/144 (61.8%)	111/133 (83.5%)	51/56 (91.1%)	<0.001 ^b
Death, %	29/333 (8.7%)	0	0	29/56 (52.8%)	< 0.001 ^b

Data are presented as a number and percentage or mean±SD or median (25th–75th percentiles); eosinophils are presented as median (fifth–95th percentiles). SPO₂, pulse oxygen saturation; ALT, alanine transaminase; AST, aspartate transaminase; HsTnI, high-sensitivity troponin; NT-proBNP, pro-brain natriuretic protein, N-terminal; SCR, serum creatinine.

^aKruskal–Wallis test.

^bWilcoxon rank-sum test.

AKI accounted for 18.2% (four of 22), but no prerenal AKI occurred. Suspected intrinsic AKI that accounted for 81.8% (18 of 22) was the most frequent form. Stage 2 comprised seven (31.8%) of 22 patients with AKI, and 50% (11 of 22) reached stage 3. The patients with AKI identified by the expanded criteria had more AKI occurrence on admission and more AKI in stage 1 than patients with AKI who met KDIGO criteria (Table 2).

Several clinical parameters were identified as being associated with proteinuria, hematuria, and AKI in patients with COVID-19 pneumonia at the time of admission, which were shown in Supplemental Tables 2– 4. Among 198 patients, one patient with AKI had a urine albumin-creatinine ratio of 238.7 μ g/mg, and his urine protein electrophoresis showed a high proportion of the renal tubular protein (40.2%).

Renal Prognosis and Risk Factors

The clinical characteristics and treatment of the 198 patients followed up on are shown in Supplemental Table 4. The mean of age was 57.1 years, 57.1%

Table 2.	Characteristics of	patients with AKI according to two different cr	riteria

22 /22 (13.6%) 9/22 (86.4%) 0/22 (0%) /22 (18.2%) 3/22 (81.8%) /22 (18.2%)	35 13/35 (37.1%) 22/35 (62.9%) 2/35 (5.7%) 4/35 (11.4%) 29/35 (82.9%)
9/22 (86.4%) 0/22 (0%) /22 (18.2%) 3/22 (81.8%) /22 (18.2%)	22/35 (62.9%) 2/35 (5.7%) 4/35 (11.4%)
0/22 (0%) /22 (18.2%) 3/22 (18.8%) /22 (18.2%)	2/35 (5.7%) 4/35 (11.4%)
/22 (18.2%) 3/22 (81.8%) /22 (18.2%)	4/35 (11.4%)
/22 (18.2%) 3/22 (81.8%) /22 (18.2%)	4/35 (11.4%)
3/22 (81.8%) /22 (18.2%)	
/22 (18.2%)	29/35 (82.9%)
	16/35 (45.7%)
/22 (31.8%)	8/35 (22.9%)
1/22 (50.0%)	11/35 (31.4%)
/22 (18.2%)	16/35 (45.7%)
_	2/2 (100%)
1/4 (25%)	1/4 (25%)
/18 (16.7%)	13/29 (44.8%)
1/4 (25.0%)	12/16 (75.0%)
3/7 (42.9%)	4/8 (50.0%)
0/11 (0%)	0/11 (0%)
6 (5–8) ^a	6 (5–11)ª
9/22 (86.4%)	20/35 (57.1%)
0/0 (0%)	0/2 (0%)
3/4 (75.0%)	3/4 (75.0%)
5/18 (88 9%)	17/29 (58.6%)
	4/16 (25.0%)
· · ·	6/8 (75.0%)
	10/11 (90.9%)
	6 (5–8) ^a 2/22 (86.4%) 0/0 (0%)

Data are presented as number and percentage. ^aThe unit of mean time for AKI recovery is day.

(113 of 198) patients were men, and the basic serum creatinine level was 74 µmol/L. A total of 98 (49.5%) of 198 patients presented with severe pneumonia and 37 (18.7%) of 198 had critical illness. Most patients received antibacterial therapy (84.8%, including moxifloxacin, levofloxacin, and cefoperazone-sulbactam), and many received antiviral therapy (umifenovir, 70.2%; lopinavir/ritonavir, 23.2%; ribavirin, 1.0%; remdesivir, 5.1%) and glucocorticoid therapy (71.7%, including at least one dose of dexamethasone in 5 mg or methylprednisolone in 20 mg, respectively).

The 198 patients were followed up on for a median duration of 12 days (Supplemental Table 5), during which 59.6% (118 of 198) patients with COVID-19 experienced pneumonia remission. Urine dipstick in 111 (68.5%) of 162 patients with proteinuria and in 44 (43.1%) of 102 patients with hematuria were reported as negative after follow-up (Table 3). According to KDIGO criteria, four (18.2%) of 22 patients with AKI achieved complete recovery of kidney function during the observation, including one of four patients with rhabdomyolysis-induced AKI, and three of 18 patients with suspected intrinsic AKI (Table 2). The mean time for AKI recovery was 6 days. The patients with AKI identified by the expanded criteria had higher rate of AKI recovery (45.7%, 16 of 35) and lower in-hospital mortality (57.1% versus 86.4%) than patients with AKI who met KDIGO criteria (Table 2). According to expanded criteria, patients with nonrecovered AKI presented with higher incidence of critical illness and severe pneumonia, and severe AKI (78.0% versus 25.0% in stage 2–3), compared with that in patients with recovered AKI (Table 4).

Stepwise multivariate binary logistic regressions were used to select significant predictors and estimate its ORs. Specifically, it showed that, compared with patients who achieved a decrease of CRP of >10 mg/L, patients whose decrease of CRP was $\leq 10 \text{ mg/L}$ had a 9.20-fold odds (95% CI, 1.62 to 52.41) of proteinuric remission (Supplemental Table 6). Moderate versus critically ill, and severe versus critically ill groups had a 4.08-fold odds (95% CI, 0.58 to 28.55) and 41.18-fold odds (95% CI, 5.36 to 316.44) of proteinuric remission, respectively. Old age (≥ 60 years) and ACEI/ARB treatment before admission were negative risk factors in proteinuric remission (old age: OR, 0.12 [95% CI, 0.02 to 0.70; ACEI/ARB: OR, 0.08 [95% CI, 0.01 to 0.68]) (Supplemental Table 6). Old age (≥ 60 years), AKI, and ACEI/ARB treatment before admission were also negative risk factors in hematuric remission (old age: OR, 0.28 [95% CI, 0.08 to 0.90]; AKI: OR, 0.12 [95% CI, 0.02 to 0.81]; ACEI/ARB: OR, 0.11 [95% CI, 0.02 to 0.73]) (Supplemental Table 7). Moderate versus critically ill, and severe versus critically ill had a 1.05-fold

RAPID COMMUNICATION www.jasn.org

Table 3.	Change in kidne	y function and biomarker	s among partici	pants in the follow-u	p study (<i>n</i> =198)

Variables	Patients before Follow-Up	Patients after Follow-Up	P Value
CRP, mg/L	56.2 (26.9–105.5)	3.9 (1.3–17.8)	< 0.001ª
Serum albumin, g/L	34.4±5.0	33.4±5.3	0.10 ^b
BUN, mmol/L	4.7 (3.5–5.9)	4.6 (3.8–6.6)	0.07 ^a
SCR, μ mol/L	74.0 (61.0–89.0)	72.5 (60.0–85.0)	0.29 ^a
Lymphocytes, 10 ⁹ /L	0.80 (0.59–1.12)	1.36 (0.87–1.82)	<0.001ª
Eosinophils, 10 ⁹ /L	0.00 (0.00-0.06)	0.00 (0.07–0.29)	<0.001ª
Serum IL-10, pg/ml	6.7 (5.0–13.1)	5.0 (5.0–5.2)	0.01 ^a
Serum IL-6, pg/ml	24.3 (12.6–57.6)	4.8 (2.0–22.4)	0.008 ^a
Serum IL-2R, U/ml	794 (552–1065)	505 (277–720)	<0.001ª
Proteinuria			<0.001ª
None	21/198 (10.6%)	132/198 (66.7%)	
±/+	146/198 (73.8%)	40/198(20.2%)	
++/+++	31/198 (15.6%)	4/198 (2.0%)	
Hematuria			<0.001ª
None	89/198 (44.9%)	133/198 (67.2%)	
±/+	84/198 (42.4%)	33/198 (16.7%)	
++/+++	25/198 (12.6%)	10/198 (5.1%)	
AKI			<0.001ª
No	163/198 (82.3%)	179/198 (90.4%)	
AKI stage 1	16/198 (8.1%)	6/198 (3.0%)	
AKI stage 2	8/198 (4.0%)	4/198 (2.0%)	
AKI stage 3	11/198 (5.6%)	9/198 (4.5%)	
Resolution on lung involvement		118/198 (59.6%)	
Death		29/198 (14.6%)	
Death in critically ill group		29/56 (51.8%)	

Data are presented as mean \pm SD or median (25th–75th percentiles) or as number and percentage. SCR, serum creatinine; $\pm/+$, \pm approximately 1+; ++/+++, 2+ approximately 3+.

^aWilcoxon signed-rank test.

^bPaired t test.

odds (95% CI, 0.21 to 5.39) and 9.16fold odds (95% CI, 1.53 to 54.97) of hematuric remission, respectively (Supplemental Table 7). Critical illness was the independently negative risk factor in the recovery of AKI in the multivariate analysis (OR, 0.03; 95% CI, 0.004 to 0.32) (Supplemental Table 8).

In the other 135 patients, 82 presented with no renal involvement at admission. The remaining 53 patients were lost to follow-up because they refused to or did not recheck urine analysis during follow-up. We compared the characteristics of the 53 patients lost to follow-up and 163 patients without AKI with follow-up. It showed no significant difference existing between the two groups (Supplemental Table 9). There was also no significant difference between the un-reported patients in this cohort and the overlapped patients in a previous cohort¹² (Supplemental Table 10).

Mortality

In this cohort study, 29 patients died during the observation period, out of 333 patients. All deaths occurred in patients with critical illness (Table 3). The patients with renal involvement, including hematuria, proteinuria, and AKI, had higher overall mortality (11.2%, 28 of 251) compared with that (1.2%, one)of 82) of patients without renal involvement (Supplemental Table 11). No one with prerenal AKI died, whereas 75% (three of four) of patients with rhabdomyolysis-induced AKI died and 58.6% (17 of 29) of patients with suspected intrinsic AKI according to expanded AKI definition died (Table 2). The mortality was 25% (four of 16) in stage 1, 75% (six of eight) in stage 2, and 90.9% (ten of 11) in stage 3, respectively (Table 2). Among patients with proteinuric remission, no one died, whereas among patients without remission, nine (17.6%) of 51 died. There were no deaths among patients with hematuric

remission; however, nine (20.3%) of 44 patients without remission died. Among patients with recovery of AKI, three (18.8%) of 16 died, whereas among patients with nonrecovery of AKI or new onset of AKI, 17 (89.5%) of 19 died (Supplemental Figure 12), suggesting that short-term outcomes of renal complications are associated with mortality in COVID-19 pneumonia.

DISCUSSION

Like Middle Eastern respiratory syndrome coronavirus (MERS-CoV), all coronaviruses have high renal involvement. The prevalence of proteinuria in this cohort study is similar to that reported in patients with MERS-CoV,¹⁹ although a quantitative assessment of urinary protein excretion was not performed and patients with CKD were excluded. In the early phase (3 weeks after symptom onset) of MERS-CoV infection,

Variables	AKI Nonrecovery	AKI Recovery	P Value
N	19	16	
Days from onset	9 (7–14)	7 (5–12)	0.22ª
Age	64.0±8.1	64.5±14.7	0.89 ^b
Sex (male)	14/19 (73.7%)	8/16 (50.0%)	0.15 ^c
Hypertension, %	9/19 (47.4%)	7/16 (43.8%)	0.83 ^c
Diabetes, %	8/19 (42.1%)	6/16 (37.5%)	0.78 ^{bc}
ACEI/ARB treatment history, %	4/19 (21.1%)	5/16 (31.3%)	0.49 ^c
Glucocorticoids treatment, %	18/19 (94.7%)	13/16 (81.3%)	0.21 ^c
Antibiotic treatment, %	19/19 (100.0%)	14/16 (87.5%)	0.11 ^c
Umifenovir treatment, %	12/19 (63.2%)	8/16 (50.0%)	0.43 ^c
Intravenous immunoglobulin therapy, %	9/19 (47.4%)	9/16 (56.3%)	0.60 ^c
Continuous KRT, %	5/19 (26.3%)	1/16 (6.3%)	0.19 ^d
Reexamined serum albumin, g/L	25.9±5.9	31.3±5.9	0.01 ^b
Reexamined lymphocytes, 10 ⁹ /L	0.47 (0.32–0.85)	0.93 (0.35–1.45)	0.07ª
Reexamined eosinophils, 10 ⁹ /L	0.03 (0.00-0.13)	0.04 (0.00-0.08)	0.67ª
SCR of baseline, μ mol/L	74.0 (61.0-89.0)	72.5 (60.0-85.0)	0.81ª
COVID-19 grade			<0.001ª
Moderate, %	1/19 (5.3%)	4/16 (25%)	
Severe, %	0/19 (0.0%)	6/16 (37.5%)	
Critically ill, %	18/19 (94.7%)	6/16 (37.5%)	
Stage of AKI			0.001ª
Stage 1	4/19 (21.1%)	12/16 (75.0%)	
Stage 2	4/19 (21.1%)	4/16 (25.0)	
Stage 3	11/19 (57.9%)	0	
Classification of AKI			0.27 ^d
Prerenal AKI	0/19 (0)	2/16 (12.5%)	
Rhabdomyolysis-induced AKI	3/19 (15.8%)	1/16 (6.3%)	
Intrinsic AKI	16/19 (84.2%)	13/16 (81.3%)	
Remission of proteinuria	1/7 (14.3%)	7/10 (70.0%)	0.05 ^d
Remission of hematuria	1/7 (14.3%)	2/6 (33.3%)	0.56 ^d

Data are presented as number and percentage, mean±SD, or median (25th-75th percentiles). SCR, serum creatinine.

^aWilcoxon rank-sum test.

^bt test.

^cChi-square test.

^dFisher exact test.

proteinuria excretion is related to fever and systemic inflammatory status,¹⁹ which is consistent with our results that half of proteinuria cases were mitigated with the recovery of pneumonia and systemic inflammation in 3 weeks after symptom onset. Therefore, we assume that proteinuria in early phase is probably transient febrile/illness-related proteinuria.

The high frequency of renal abnormalities in this study, including 75.4% with renal involvement, 65.8% with proteinuria, and 41.7% with hematuria, were similar to findings in other patients with critical illness, in which 67.4% had proteinuria and 77.5% had hematuria,²⁰ suggesting proteinuria and hematuria in COVID-19 is not different from patients with other critical illnesses. However, the incidence of AKI is lower.

About 7.5% (35 of 467) of the patients in the overall cohort had AKI according to the expanded criteria; and 45.7% of these patients had stage 1 AKI and 45.7% (16 of 35) achieved complete remission in 3 weeks after onset of disease. The mean time for AKI recovery was 6 days. When compared with patients with AKI who met KDIGO criteria, although the patients with AKI identified by the expanded criteria presented with more AKI on admission, they had a higher rate of AKI recovery and lower in-hospital mortality. As expanded criteria helped to identify the recovery of AKI, these data suggested that half of the early AKI in patients with COVID-19 was mild and easily recovered.

Using either KDIGO criteria or expanded criteria, suspected intrinsic AKI

accounted for the most frequent form of AKI (>80%). Pathology from autopsies of patients with COVID-19 with renal function impairment demonstrated the direct evidence of intrinsic renal involvement, and revealed that the kidneys had varying degrees of ATN, luminal brush border sloughing, hyaline casts, microthrombi, and mild fibrosis in the interstitium, whereas severe glomerular injury and lymphocyte infiltration were absent,10,21 suggesting the major form of intrarenal AKI in COVID-19 is ATN, which is consistent with the high proportion of renal tubular protein in urine findings. The high rate of resolution of proteinuria, hematuria, and AKI in the early stage (3 weeks after onset of symptom) is also consistent with the postmortem data that ATN appears to be the most common cause. However, the mortality in patients with renal involvement, especially in those with no improvement of kidney function during the follow-up period (89.5%, 17 of 19), was extremely high in the early stage, suggesting that despite good short-term renal prognosis, renal complications in COVID-19 still remain associated with poor mortality.

Risk factors and causes of AKI in COVID-19 are diverse and multifactorial. On the basis of our multivariate binary regression analysis, we suggest that severity of pneumonia is the most important factor in development of AKI in patients with COVID-19. The fundamental pathophysiology of pneumonia in critically ill patients is severe acute respiratory distress syndrome, which has been identified as an independent risk factor associated with occurrence of AKI.²² These data suggest it is necessary for intensively supporting and carefully monitoring patients with severe and critically ill pneumonia to ameliorate renal complications.

This study also has several limitations. First, the number of patients included in this study is limited, and there were some missing data. Second, because of the strain on medical resources in the epicenter of the COVID-19 outbreak, we could not get full laboratory support to obtain more sufficient evidence for evaluating the urine. Third, the duration of observation is not long enough to implement survival analysis to predict the probabilities for remission of renal damage in the long term and risks for mortality. Fourth, we did not detect SARS-CoV-2 in urine samples. Therefore, we could not assess correlations between urine virus and renal complications. Fifth, we did not have renal pathology data to assess the direct effect of the virus on renal outcomes. We did not detect biomarkers, such as urinary NGAL and KIM-1, to accurately distinguish prerenal AKI from ATN. Sixth, as the admission creatinine was used to define AKI in expanded criteria, some AKI at admission that did not resolve may have not been included, which might potentially overestimate overall

recovery as well as underestimate severity in some patients (those with AKI at admission that continued to worsen).

In conclusion, despite high morbidity of renal involvement, the short-term renal prognosis of patients is good, as half of them achieved remission in 3 weeks after onset of symptoms. However, adverse short-term outcomes of renal involvement are also associated with mortality in COVID-19. Severity of pneumonia was identified as an independent negative prognostic indicator for renal complications. Therefore, the strategy on treatment and prevention of severe or critically ill pneumonia is appropriate for COVID-19–related renal complications.

ACKNOWLEDGMENTS

The authors greatly appreciate all of the hospital staff for their efforts in recruiting and treating patients and thank all patients involved in this study.

Dr. Xu, Dr. Yao, and Dr. Zeng designed the study. Dr. Pei, Dr. Liu, Dr. Zhang, Dr. Yu, Dr. Ma, Dr. Huang, and Dr. Liu collected the data. Dr. Pei, Dr. Zhang, and Dr. Zeng prepared figures and tables. Dr. Pei and Dr. Zhang contributed analytical tools. Dr. Pei, Dr. Zhang, and Dr. Zeng wrote the paper. Dr. Zeng and Dr. Yao conceived the project and supervised and coordinated all of the work.

DISCLOSURE

All authors declare that they have no competing interests, no support from any organization for the submitted work, and no financial relationships with any organizations that might have an interest in the submitted work.

FUNDING

This work was financially supported by international (regional) cooperation and exchange projects (National Natural Science Foundation of China-Deutsche Forschungsgemeinschaft, grant 81761138041), the Major Research Plan of the National Natural Science Foundation of China (grant 91742204), the National Natural Science Foundation of China (grants 81670633, 81600556, 81770681, and 81974086), and the National Key Research and Development Program (grants 2016YFC0906103 and 2018YFC1314000).

SUPPLEMENTAL MATERIAL

This article contains the following supplemental material online at http://jasn. asnjournals.org/lookup/suppl/doi:10.1681/ ASN.2020030276/-/DCSupplemental.

Supplemental Table 1. Hematuria and proteinuria in the non-AKI group versus the AKI group.

Supplemental Table 2. The correlation between proteinuria and variables.

Supplemental Table 3. The correlation between hematuria and variables.

Supplemental Table 4. The correlation between AKI occurrence and variables.

Supplemental Table 5. Baseline and treatment characteristics among participants in the follow-up study (N=198).

Supplemental Table 6. Risk factors associated with proteinuria remission during follow-up periods.

Supplemental Table 7. Risk factors associated with hematuria remission during follow-up periods.

Supplemental Table 8. Risk factors associated with AKI recovery during follow-up periods.

Supplemental Table 9. Characteristics of patients lost to follow-up and patients with follow-up.

Supplemental Table 10. Mortality of the patients with renal involvement during follow-up periods.

Supplemental Table 11. Comparison of unreported patients in this cohort and the overlapped patients in a previous cohort.

Supplemental Figure 12. Relationship between outcomes of renal involvement and mortality during follow-up periods.

REFERENCES

- Li Q, Guan X, Wu P, Wang X, Zhou L, Tong Y, et al.: Early transmission dynamics in Wuhan, China, of novel coronavirus-infected pneumonia. N Engl J Med 382: 1199–1207, 2020
- Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, et al.; China Novel Coronavirus Investigating and Research Team: A novel coronavirus from patients with pneumonia in

China, 2019. N Engl J Med 382: 727–733, 2020

- Chen N, Zhou M, Dong X, Qu J, Gong F, Han Y, et al.: Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: A descriptive study. *Lancet* 395: 507–513, 2020
- Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al.: Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China [published online ahead of print Feb 7, 2020]. JAMA 10.1001/ jama.2020.1585
- Smith N, Fraser M: Straining the system: Novel coronavirus (COVID-19) and preparedness for concomitant disasters. Am J Public Health 110: 648–649, 2020
- Velavan TP, Meyer CG: The COVID-19 epidemic. Trop Med Int Health 25: 278–280, 2020
- Ki M; Task Force for 2019-nCoV: Epidemiologic characteristics of early cases with 2019 novel coronavirus (2019-nCoV) disease in Korea. Epidemiol Health 42: e2020007, 2020
- Lu R, Zhao X, Li J, Niu P, Yang B, Wu H, et al.: Genomic characterisation and epidemiology of 2019 novel coronavirus: Implications for virus origins and receptor binding. *Lancet* 395: 565–574, 2020
- Chan JF, Yuan S, Kok KH, To KK, Chu H, Yang J, et al.: A familial cluster of pneumonia associated with the 2019 novel coronavirus indicating person-to-person transmission: A study of a family cluster. *Lancet* 395: 514–523, 2020

- National Health Commission of the People's Republic of China: Diagnosis and Treatment Protocols of Pneumonia caused by a Novel Coronavirus (Trial Version 7), 2020. Available at: http://www.nhc.gov.cn/yzygj/s7653p/202003/ 46c9294a7dfe4cef80dc7f5912eb1989/files/ ce3e6945832a438eaae415350a8ce964.pdf. Accessed March 3, 2020
- Pan F, Ye T, Sun P, Gui S, Liang B, Li L, et al.: Time course of lung changes on chest CT during recovery from 2019 novel coronavirus (COVID-19) pneumonia [published online ahead of print Feb 13, 2020]. Radiology 10.1148/radiol.2020200370
- Cheng Y, Luo R, Wang K, Zhang M, Wang Z, Dong L, et al.: Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int* 5: 829–838, 2020 10.1016/j.kint.2020.03.005
- Yang L, Xing G, Wang L, Wu Y, Li S, Xu G, et al.; ISN AKF 0by25 China Consortiums: Acute kidney injury in China: A crosssectional survey. *Lancet* 386: 1465–1471, 2015
- Khwaja A: KDIGO clinical practice guidelines for acute kidney injury. Nephron Clin Pract 120: c179–c184, 2012
- Parikh CR, Coca SG: Acute kidney injury: Defining prerenal azotemia in clinical practice and research. Nat Rev Nephrol 6: 641–642, 2010
- Lim YS, Cho H, Lee ST, Lee Y: Acute kidney injury in pediatric patients with rhabdomyolysis. Korean J Pediatr 61: 95–100, 2018

- Stahl K, Rastelli E, Schoser B: A systematic review on the definition of rhabdomyolysis. *J Neurol* 267: 877–882, 2020
- Kidney Disease: Improving Global Outcomes (KDIGO) CKD Work Group: KDIGO 2012 clinical practice guideline for the evaluation and management of chronic kidney disease. *Kidney Int Suppl* 3: 1–150, 2013
- Cha RH, Joh JS, Jeong I, Lee JY, Shin HS, Kim G, et al.; Critical Care Team of National Medical Center: Renal complications and their prognosis in Korean patients with Middle East respiratory syndrome-coronavirus from the central MERS-CoV designated hospital. J Korean Med Sci 30: 1807–1814, 2015
- Han SS, Ahn SY, Ryu J, Baek SH, Chin HJ, Na KY, et al.: Proteinuria and hematuria are associated with acute kidney injury and mortality in critically ill patients: A retrospective observational study. *BMC Nephrol* 15: 93, 2014
- Diao B, Wang C, Wang R, Feng Z, Tan Y, Wang H, et al.: Human kidney is a target for novel severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. medRxiv p. 2020.03.04.20031120: 2020
- Darmon M, Clec'h C, Adrie C, Argaud L, Allaouchiche B, Azoulay E, et al.: Acute respiratory distress syndrome and risk of AKI among critically ill patients. *Clin J Am Soc Nephrol* 9: 1347–1353, 2014